



Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to MDH for the week ending January 27, 2018

Prepared by the Infectious Disease Epidemiology and Outbreak Response Bureau
Prevention and Health Promotion Administration
Maryland Department of Health

The data presented in this document are provisional and subject to change as additional reports are received.

SUMMARY

During the week ending January 27, 2018, influenza-like illness (ILI) intensity in Maryland was **HIGH** and there was **WIDESPREAD** geographic activity. The proportion of outpatient visits for ILI reported by Sentinel Providers and by Maryland Emergency Departments increased. The proportion of MRITS respondents reporting ILI decreased. Clinical laboratories reported an increase in the proportion of specimens testing positive for influenza. One hundred thirty-four specimens tested positive for influenza at the MDH lab. There were 314 influenza-associated hospitalizations. Twenty-five respiratory outbreaks were reported to MDH.


[Click here to visit our influenza surveillance web page](#)

ILI Intensity Levels
Minimal
Low
Moderate
✓ High

Influenza Geographic Activity
No Activity
Sporadic
Local
Regional
✓ Widespread

ILINet Sentinel Providers

Twenty-five sentinel providers reported a total of 5,898 visits this week. Of those, 404 (6.8%) were visits for ILI. This is **above** the Maryland baseline of 2.0%.



ILI Visits To Sentinel Providers By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	83 (21%)	87 (25%)	670 (26%)
Age 5-24	118 (29%)	119 (35%)	918 (35%)
Age 25-49	113 (28%)	75 (22%)	577 (22%)
Age 50-64	66 (16%)	37 (11%)	289 (11%)
Age ≥ 65	24 (6%)	26 (8%)	152 (6%)
Total	404 (100%)	344 (100%)	2,606 (100%)

Visits to Emergency Departments for ILI

Emergency Departments in Maryland reported a total of 45,427 visits this week through the [ESSENCE surveillance system](#). Of those, 3,178 (7.0%) were visits for ILI.



ILI Visits To Emergency Departments By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	591 (19%)	466 (20%)	5,716 (28%)
Age 5-24	914 (29%)	574 (25%)	5,007 (25%)
Age 25-49	853 (27%)	682 (29%)	5,032 (25%)
Age 50-64	455 (14%)	364 (16%)	2,442 (12%)
Age ≥ 65	365 (11%)	245 (11%)	1,930 (10%)
Total	3,178 (100%)	2,331 (100%)	20,127 (100%)

Neighboring states' influenza information:


- Delaware <http://dhss.delaware.gov/dph/epi/influenzahome.html>
- District of Columbia <http://doh.dc.gov/service/influenza>
- Pennsylvania <http://www.health.pa.gov/My%20Health/Diseases%20and%20Conditions/I-L/Pages/Influenza.aspx#.V-LtaPkrJD8>
- Virginia <http://www.vdh.virginia.gov/epidemiology/influenza-flu-in-virginia/influenza-surveillance/>
- West Virginia <http://dhhr.wv.gov/oeps/disease/flu/Pages/fluSurveillance.aspx>

Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to MDH for the week ending January 27, 2018

Community-based Influenza Surveillance (MRITS)


MRITS is the Maryland Resident Influenza Tracking System, a weekly survey for influenza-like illness (ILI). A total of 670 residents responded to the [MRITS survey](#) this week. Of those, 14 (2.1%) reported having ILI and missing 31 cumulative days of regular daily activities.




MRITS Respondents Reporting ILI By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	--	1 (5%)	5 (3%)
Age 5-24	4 (29%)	3 (16%)	41 (28%)
Age 25-49	3 (21%)	8 (42%)	35 (24%)
Age 50-64	5 (36%)	6 (32%)	52 (35%)
Age ≥ 65	2 (14%)	1 (5%)	14 (10%)
Total	14 (100%)	19 (100%)	147 (100%)

Clinical Laboratory Influenza Testing

There were 58 clinical laboratories reporting 9,529 influenza diagnostic tests, mostly rapid influenza diagnostic tests (RIDTs). Of those, 3,018 (31.7%) were positive for influenza. Of those testing positive, 2,323 (77%) were influenza Type A and 695 (23%) were influenza Type B. The [reliability of RIDTs](#) depends largely on the conditions under which they are used. False-positive (and true-negative) results are more likely to occur when the disease prevalence in the community is low, which is generally at the beginning and end of the influenza season and during the summer.




Positive Rapid Flu Tests by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A	2,323 (77%)	1,929 (81%)	9,047 (80%)
Type B	695 (23%)	452 (19%)	2,247 (20%)
Total	3,018 (100%)	2,381 (100%)	11,294 (100%)



State Laboratories Administration Influenza Testing

The MDH Laboratories Administration performed a total of 224 PCR tests for influenza and 134 (59.8%) were positive for influenza. Of those testing positive, 96 (71.6%) were positive for Type A (H3), 18 (13.4%) were positive for Type A (H1), 19 (14.2%) were positive for Type B (Yamagata), and 1 (0.7%) was positive for dual Type A(H3)/B. PCR testing is more reliable than RIDT. The MDH testing identifies subtypes of influenza A and lineages of influenza B, information that is not available from the RIDT results. The table below summarizes results by type, subtype, and lineage.



Positive PCR Tests by Type (Subtype)	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A (H1)	18 (13%)	53 (22%)	198 (17%)
Type A (H3)	96 (72%)	180 (73%)	822 (72%)
Type B (Victoria)	--	--	4 (<1%)
Type B (Yamagata)	19 (14%)	12 (5%)	105 (9%)
Dual Type A (H1/H3)	--	--	1 (<1%)
Dual Type A(H3)/B	1 (1%)	1 (<1%)	2 (<1%)
Type A (H3N2v)	--	--	3 (<1%)
Total	134 (100%)	246 (100%)	1,135 (100%)

Where to get an influenza vaccination


Interested in getting a flu vaccine for the 2017-18 influenza season? Go to <https://phpa.health.maryland.gov/influenza/Pages/getvaccinated.aspx> and click on your county/city of residence. You will be redirected to your local health department website for local information on where to get your flu vaccine.

Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to MDH for the week ending January 27, 2018

Influenza-associated Hospitalizations


A total of 314 influenza-associated hospitalizations were reported this week. (A person with an overnight hospital stay along with a positive influenza test of any kind, e.g., RIDT or PCR, is considered an “influenza-associated hospitalization” for purposes of influenza surveillance.)



Influenza-Associated Hospitalizations by Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	30 (10%)	17 (5%)	115 (7%)
Age 5-17	18 (6%)	13 (4%)	68 (4%)
Age 18-24	10 (3%)	5 (2%)	45 (3%)
Age 25-49	31 (10%)	38 (12%)	214 (12%)
Age 50-64	68 (22%)	67 (21%)	367 (21%)
Age ≥ 65	157 (50%)	176 (56%)	949 (54%)
Total	314 (100%)	316 (100%)	1,758 (100%)

Outbreaks of Respiratory Disease

There were twenty-five respiratory outbreaks reported to MDH this week. (Disease outbreaks of any kind are reportable in Maryland. Respiratory outbreaks may be reclassified once a causative agent is detected, e.g., from ILI to influenza.)



Respiratory Outbreaks by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Influenza	21 (84%)	12 (67%)	57 (68%)
Influenza-like Illness	2 (8%)	4 (22%)	18 (21%)
Pneumonia	2 (8%)	2 (11%)	9 (11%)
Other Respiratory	--	--	--
Total	25 (100%)	18 (100%)	84 (100%)

National Influenza Surveillance (CDC)

During week 4 (January 21-27, 2018), influenza activity increased in the United States.

- Viral Surveillance:** The most frequently identified influenza virus subtype reported by public health laboratories during week 4 was influenza A(H3). The percentage of respiratory specimens testing positive for influenza in clinical laboratories remained elevated.
- Pneumonia and Influenza Mortality:** The proportion of deaths attributed to pneumonia and influenza (P&I) was above the system-specific epidemic threshold in the National Center for Health Statistics (NCHS) Mortality Surveillance System.
- Influenza-associated Pediatric Deaths:** Seventeen influenza-associated pediatric deaths were reported, one of which occurred during the 2015-2016 season.
- Influenza-associated Hospitalizations:** A cumulative rate of 51.4 laboratory-confirmed influenza-associated hospitalizations per 100,000 population was reported.
- Outpatient Illness Surveillance:** The proportion of outpatient visits for influenza-like illness (ILI) was 7.1%, which is above the national baseline of 2.2%. All 10 regions reported ILI at or above region-specific baseline levels. New York City, the District of Columbia, and 42 states experienced high ILI activity; Puerto Rico and two states experienced moderate ILI activity; three states experienced low ILI activity; and three states experienced minimal ILI activity.
- Geographic Spread of Influenza:** The geographic spread of influenza in Puerto Rico and 48 states was reported as widespread; Guam and one state reported regional activity; the District of Columbia and one state reported local activity; and the U.S. Virgin Islands reported sporadic activity.

